

Can a quantum measurement be cancelled in a very short period of time?

R. L. Schafr^{*}

CISM, London Metropolitan University, London EC3N 1JY, U.K.

Abstract

A test is suggested for whether the obtaining of certain information, and then deleting it too quickly to be retained, constitutes a quantum measurement of that information.

The question of what exactly constitutes a measurement, and when it takes place, remains debatable after more than 75 years of quantum mechanics. Here a method will be suggested for testing whether the capture of certain information by macroscopic instruments, followed by its partial erasure after a very short period of time, constitutes a measurement of all the information that was first captured, or only of the information which remains after the erasure.

Consider a system of two 2-level particles, for instance two spin- $\frac{1}{2}$ particles, and suppose them in an entangled state. Then if the particle spins are measured in directions θ and ϕ respectively, this is a measurement of the two commuting observables

$$(\sigma^A_{\theta} \otimes id, id \otimes \sigma^B_{\phi}) \quad (1)$$

and we can deduce, in the usual way, that measurement of even one of these observables projects both particles into two definite unentangled states, and measurements can be made which will confirm, for instance, the correlations between them.

Suppose instead that we measured

$$\sigma^A_{\theta} \otimes \sigma^B_{\phi} \quad (2)$$

This observable represents (as it indicates) the product of the individual particles' outcomes, and, if the outcomes are treated as ± 1 , can be regarded as measuring whether their results are the same (eigenvalue $+1$) or opposite (eigenvalue -1).

If a measurement is made, choosing either the observables (1) or the observable (2), the results are compatible, in that the probabilities of getting the results "same" or "opposite" are the same. But if two successive such measurements are made at different angles, incompatible results can be obtained¹. For example, suppose the two particles are initially in the (normalized) state

$$\sum_{i,j=\pm 1} \alpha_{ij} |i,j\rangle \quad (3)$$

^{*}e-mail: roger.schafr@londonmet.ac.uk

¹This was brought to my attention by G. Adenier [1], though Adenier argued for a viewpoint which is not adopted here.

with respect to z-direction eigenstates, and that, first, both are measured in the z-direction, and then, second, a measurement is made of the observables in (1) using the x-direction. If the first measurement is of the observables (1), the system is projected into one of the states $|i, j\rangle$ with probabilities $|\alpha_{ij}|^2$, and if both particles are then measured in the x-direction, each x-eigenstate is obtained with equal probability, and so in an ensemble of the two successive measurements there will be equal probability for each of the four outcomes at the end of the second measurement. But if the first measurement (in the z-direction) is of the observable in (2), then the system is projected into one of the states

$$\frac{1}{\sqrt{p_1}} \{ \alpha_{11} |1, 1\rangle + \alpha_{-1-1} |-1, -1\rangle \}, \quad \frac{1}{\sqrt{p_{-1}}} \{ \alpha_{1-1} |1, -1\rangle + \alpha_{-11} |-1, 1\rangle \} \quad (4)$$

where p_1, p_2 are the respective probabilities, given by

$$p_1 = |\alpha_{11}|^2 + |\alpha_{-1-1}|^2, \quad p_{-1} = |\alpha_{1-1}|^2 + |\alpha_{-11}|^2 \quad (5)$$

and then the probability of, for instance, obtaining the x-eigenstate $|1, 1\rangle_x$ at the end of the second measurement is

$$\frac{1}{4} \{ |\alpha_{11} + \alpha_{-1-1}|^2 + |\alpha_{1-1} + \alpha_{-11}|^2 \} \quad (6)$$

which is clearly $\neq \frac{1}{4}$ (except for special values of the α_{ij}). It is also generally different to the probability of $|1, 1\rangle_x$ if the first measurement is omitted, which is

$$\frac{1}{4} \{ |\alpha_{11} + \alpha_{-1-1} + \alpha_{1-1} + \alpha_{-11}|^2 \} \quad (7)$$

If we consider how this could be verified by experiment, the obvious question is how to measure whether the observers' results are the same or opposite while leaving the individual results themselves unknown, and indeed unknowable.² But this question can be turned into a test of quantum measurement. Suppose the particles are brought together (rather than being measured while spacelike-separated), and the measurement is made by an apparatus which is considered macroscopic, and which measures the two outcomes, multiplies them and retains that result, then irretrievably deletes the original individual results. If more than a very short time elapses between making the individual measurements and deleting them, the results must be appropriate to a measurement of (1), since the individual results could have been noted and recorded (whether they actually were or not). But if deletion takes place in a time so short that it was impossible to record the original results, then we are in the area of ignorance about what exactly constitutes a quantum measurement, and it is conjectural whether the results would be appropriate to (1) or to (2).

The situation can obviously be generalised, and does not necessarily need entangled states. Suppose $\hat{A}$ and $\hat{B}$ are a complete set of commuting observables on a system, i.e. $\hat{A}$ and $\hat{B}$ are degenerate but the degeneracy is lifted when both are measured. Suppose the joint measurement is made, and then, very quickly, a function $f(\hat{A}, \hat{B})$ is evaluated from the outcomes, and recorded by methods which would usually be regarded as classical, while the values of $\hat{A}$ and $\hat{B}$ themselves are discarded

²But see in this context Ref [2]

irreversibly. Provided $f(\hat{A}, \hat{B})$ is also degenerate, the projection by $f(\hat{A}, \hat{B})$ is not generally the same as projection by the pair $(\hat{A}, \hat{B})$. So would subsequent measurements give the results expected if the first measurement was of $(\hat{A}, \hat{B})$, or $f(\hat{A}, \hat{B})$?

This article will not attempt to design a specific experimental apparatus, but the requirements of an apparatus are as follows. We want the calculation and recording of the value of $f(\hat{A}, \hat{B})$ from $(\hat{A}, \hat{B})$ to take place by classical means, and we want the subsequent deletion of $(\hat{A}, \hat{B})$ to be classical, yet irreversible. So suppose that the multiplication is carried out by some standard type of computer or calculator, while the deletion of the original results takes place before any signal could reach the memory section of the apparatus, and/or could be retained by the memory in view of the known volatility and other properties of the materials used. Of course it might be that a different design of apparatus and different materials could change the time threshold for the deletion, but that would be a different experiment; what matters is that with the particular design of apparatus which has been chosen, the information has been discarded irretrievably before it could be recorded, even if the apparatus was working at maximum efficiency.

If there is considered to be a difficulty in measuring a degenerate observable without lifting the degeneracy by co-measuring a compatible observable (at least accidentally), we may regard the measurement of $f(\hat{A}, \hat{B})$ as part of a non-degenerate measurement one of whose outcomes in each degeneracy eigenspace, and the only one which can be obtained from the initial state since the others are orthogonal to it, is the same state as a direct application of the projector into the degeneracy eigenspace. For instance in the entangled particles example, if the outcomes of projection by $(\sigma^A_z \otimes \sigma^B_z)$ which are given in (4) are supplemented by orthogonal states, i.e. by

$$\frac{1}{\sqrt{p_1}} \{-\alpha_{-1-1} |1, 1\rangle + \alpha_{11} |-1, -1\rangle\}, \quad \frac{1}{\sqrt{p_{-1}}} \{-\alpha_{-11} |1, -1\rangle + \alpha_{1-1} |-1, 1\rangle\} \quad (8)$$

we have a basis for a degeneracy-lifting co-measurement of $(\sigma^A_z \otimes \sigma^B_z)$ which gives the same outcomes, with the same probabilities, as a measurement of $(\sigma^A_z \otimes \sigma^B_z)$ on its own.

If this experiment, or more generally an experiment for $f(\hat{A}, \hat{B})$ versus $(\hat{A}, \hat{B})$, showed that the second measurement gave results appropriate to the first measurement being of $f(\hat{A}, \hat{B})$, despite the fact that the only way $f(\hat{A}, \hat{B})$ could have been evaluated was from the lost results of $(\hat{A}, \hat{B})$ (nor could the recorded value of $f(\hat{A}, \hat{B})$ be simply a mistake, with no first measurement having really happened in the circumstances), it must mean that what was measured was somehow relative to events at the macroscopic level. This would be compatible with the Copenhagen view that a quantum situation resides partly in the macroscopic context, and is also consistent with the “irreversible record” viewpoint that a measurement takes place when a record is created by an irreversible process. If the second measurement showed that the first measurement was of $(\hat{A}, \hat{B})$, this would be consistent with “realist” pictures, decoherence, and also the many-worlds interpretation, since a splitting of observers should take place when $(\hat{A}, \hat{B})$ is measured, with no splitting for the classical process of evaluating $f(\hat{A}, \hat{B})$ and discarding $(\hat{A}, \hat{B})$. Which of the two possibilities would be found seems to be an open question.

References

- [1] G. Adenier, quant-ph/0105031
- [2] C. Simon, M.Zukowski, H. Weinfurter and A. Zeilinger, Phys. Rev. Lett. **85**, 1783 (2000)(quant-ph/0009074); R.L. Schafir, quant-ph/0208127